
MYCOTAXON

<http://dx.doi.org/10.5248/117.429>

Volume 117, pp. 429–434

July–September 2011

New recombinations in *Glomeromycota*

FRITZ OEHL^{1*}, GLADSTONE ALVES DA SILVA²,
BRUNO TOMIO GOTO³ & EWALD SIEVERDING⁴

¹Federal Research Institute Agroscope Reckenholz-Tänikon ART, Organic Farming Systems,
Reckenholzstrasse 191, CH-8046 Zürich, Switzerland

²Departamento de Micologia, CCB, Universidade Federal de Pernambuco,
Av. Prof. Nelson Chaves s/n, Cidade Universitária, 50670-420, Recife, PE, Brazil

³Departamento de Botânica, Ecologia e Zoologia, CB, Universidade Federal do Rio Grande
do Norte, Campus Universitário, 59072-970, Natal, RN, Brazil

⁴Institute for Plant Production and Agroecology in the Tropics and Subtropics,
University of Hohenheim, Garbenstrasse 13, D-70599 Stuttgart, Germany

*CORRESPONDENCE TO: fritz.oehl@art.admin.ch

ABSTRACT — Combined morphological and molecular biological analyses have achieved major advances in the taxonomy of fungal species in phylum *Glomeromycota*. In this study, we analyzed which species might not yet be attributed to their correct genus, focusing on *Acaulospora myriocarpa*, *A. undulata*, *A. nicolsonii*, and *Scutellospora nodosa*. Based on spore wall structure and phylogenetic support, we recombined the taxa as *Archaeospora myriocarpa*, *Ar. undulata*, *Ambispora nicolsonii*, and *Cetraspora nodosa*.

KEY WORDS — molecular phylogeny, rDNA

Introduction

Combined morphological and molecular phylogenetic analyses have thoroughly changed the classification of the phylum *Glomeromycota* (e.g. Morton & Redecker 2001, Oehl et al. 2008, 2011a). Today, there are three glomeromycotan classes, five orders, 14 families, and 29 genera (Oehl et al. 2011a,d). When species were attributed to these newly established higher taxa, it was known that some species were incorrectly placed due to insufficient available morphological or phylogenetical information (Oehl et al. 2008, 2011a,b,e). In this paper, we consider four of those species — *Acaulospora myriocarpa*, *Ac. undulata*, *Ac. nicolsonii*, and *Scutellospora nodosa*. As a result of new available information, we transfer these species to their proper genera based on their morphological features.

Material & methods

Specimens

Holotype or isotype materials of *Ac. myriocarpa* (OSC, herbarium of Oregon State University, Corvallis, U.S.A.), *Ac. undulata* (OSC), and *Sc. nodosa* (DPP, the Plant Pathology herbarium, Academy of Agriculture, Szczecin, Poland) were analyzed. Type material of *Ac. nicolsonii* (Walker et al. 1984) was either never deposited at OSC or lost during shipping (Richard Halse & Joey Spatafora, curators of OSC, pers. communication); therefore here we rely on the reinterpretation by Walker et al. (1984) of the original species description and illustrations and on comparison of *Ac. nicolsonii* with the type materials of *Ambispora gerdemannii* and *Am. fennica* in Walker et al. (2007).

Morphological analyses

Morphological descriptions of spores (including sporiferous saccules, sporogenous cells, germination shields, and subcellular structures) are based on observations of specimens mounted in polyvinyl alcohol-lactic acid-glycerol (PVLG) and in a mixture of PVLG+Melzer's reagent (Brundrett et al. 1994). Spore structure terminology follows Goto & Maia (2006), Sieverding & Oehl (2006), Oehl et al. (2006), Spain et al. (2006), Palenzuela et al. (2008, 2010, 2011), Goto et al. (2008) and Oehl et al. (2011a–c) for acaulosporoid, ambisporoid, archaeosporoid, entrophosporoid, and gigasporoid spore forming species.

Results

Acaulospora myriocarpa and *Acaulospora undulata*

Both *Ac. myriocarpa* and *Ac. undulata* form bi-walled spores laterally on the neck of a sporiferous saccule. The outer wall is bi-layered and the inner wall triple-layered. The staining reaction of the mycorrhizal structures is faint to absent, and vesicle formation has been observed only infrequently in pure cultures of the two species (Schenck et al. 1986) and then possibly due to misinterpretation (Sieverding own observations). These features attribute both species unequivocally to the genus *Archaeospora* J.B. Morton & D. Redecker emend. Spain (2003).

Archaeospora myriocarpa (Spain, Sieverd. & N.C. Schenck) Oehl, G.A. Silva, B.T. Goto & Sieverd., **comb. nov.**

MYCOBANK MB 561657

= *Acaulospora myriocarpa* Spain, Sieverd. & N.C. Schenck, Mycotaxon 25: 112. 1986.

Archaeospora undulata (Sieverd.) Sieverd., G.A. Silva, B.T. Goto & Oehl, **comb. nov.**

MYCOBANK MB 561662

= *Acaulospora undulata* Sieverd. Angew. Bot. 62: 373. 1988.

Acaulospora nicolsonii

Acaulospora nicolsonii regularly forms spores on the lateral branch (i.e. pedicel) of a sporiferous saccule. Its spores have three walls: an outer wall, which during degradation shows cracking fissures known only for *Ambispora*

species, a middle wall, and an inner wall. Since the spore morphology exactly matches the features of typical acaulosporoid spores in *Ambispora* C. Walker et al. (= 'acaulo-ambisporoid' spores), we transfer the species to this genus (Spain et al. 2006, Walker et al. 2007, 2008).

Ambispora nicolsonii (C. Walker, L.E. Reed & F.E. Sanders) Oehl, G.A. Silva, B.T. Goto & Sieverd., **comb. nov.**

MYCOBANK MB 561663

= *Acaulospora nicolsonii* C. Walker, L.E. Reed & F.E. Sanders,
Trans Br. Mycol. Soc. 83: 360. 1984.

Scutellospora nodosa

New observations on the germination shield of the type material of *Sc. nodosa* reveal that the shields are hyaline and multiply lobed, which is typical for *Racocetraceae* species (e.g. Oehl et al. 2010). Since the species has triple-walled spores (Błaszczkowski 1991, Oehl et al. 2008), we transfer it to the genus *Cetraspora* Oehl et al. (*Racocetraceae*).

Cetraspora nodosa (Błaszcz.) Oehl, G.A. Silva, B.T. Goto & Sieverd., **comb. nov.**

MYCOBANK MB 561664

= *Scutellospora nodosa* Błaszcz., Mycologia 83: 537. 1991.

Discussion

Archaeospora has now three species— two with smooth spore walls (*Ar. trappei*, *Ar. myriocarpa*) and one with an ornamented (undulating) spore surface (*Ar. undulata*).

Ambispora has now three species (*Am. callosa*, *Am. fecundispora*, *Am. leptoticha*) for which only glomoid spore formation is known and two species (*Am. jimgerdemannii*, *Am. nicolsonii*) for which only acaulosporoid spore formation is known. All other *Ambispora* species have spore bimorphism (*Am. appendicula*, *Am. brasiliensis*, *Am. fennica*, *Am. gerdemannii*, *Am. granatensis*) (Spain et al. 2006, Walker 2008, Goto et al. 2008, Palenzuela et al. 2011).

Three major characteristics strongly support transfer of *Ac. nicolsonii* to *Ambispora*: i) the presence of three walls (outer, middle, inner), ii) the clear pedicel on the sporiferous saccule neck, referred to as a 'stalk' in the protologue (Walker et al. 1984), and iii) outer spore walls that have the typical 'cracking fissures' as known for outer walls of almost all known *Ambispora* species, e.g. *Am. appendicula*, *Am. gerdemannii*, *Am. fennica*, and *Am. brasiliensis* (see illustrations in Walker et al. 1984; Spain et al. 2006, Goto et al. 2008). The absence of fracturing (i.e., 'plate-like splitting') of the middle wall layers in *Am. nicolsonii* (Walker et al. 2007) might be the most important morphological difference between it and *Am. gerdemannii* and *Am. fennica*, while the outer walls in both *Am. nicolsonii* and *Am. gerdemannii* do not stain in Melzer's reagent but do in *Am. fennica* (Walker et al. 2007; Oehl, pers. observation).

The recent transfer of *Am. brasiliensis* to *Acaulospora* by Krüger et al. (2011) cannot be accepted; the spore morphology of the Scottish material described and illustrated as *Acaulospora brasiliensis* by Krüger et al. (2011) does not match the morphology of *Am. brasiliensis*, and their sequenced material apparently represents a true *Acaulospora* species rather than authentic *Am. brasiliensis*. Krüger et al. (2011) unfortunately ignored the major characters elaborated by Spain et al. (2006) for *Ambispora* species and detailed by Goto et al. (2008) for *Am. brasiliensis*.

There have been doubts about the shield morphology of *Cetraspora nodosa* (*Scutellospora nodosa* in Oehl et al. 2008), and sequences of the fungus deposited in public databases during more than 10 years were apparently misidentified. New combined morphological and molecular phylogenetic analyses, based on identified sequences recently placed in the public databases, now show a congruency between morphological and molecular data supporting *C. nodosa* as closely related to *C. helvetica*, *C. pellucida*, and *C. gilmorei* (Oehl et al. 2010, 2011b) within the monophyletic genus *Cetraspora*.

Acknowledgments

We acknowledge the valuable comments and revisions of Felipe Wartchow (Universidade Federal de Paraíba, João Pessoa, Brazil), Eduardo Furrázola (Instituto de Ecología y Sistemática, La Habana, Cuba), and Javier Palenzuela (EEZ, CSIC, Granada, Spain) on the manuscript and appreciate the corrections by Shaun Pennycook, Nomenclature Editor, and suggestions by Lorelei L. Norvell, Editor-in-Chief. This study was supported by the Swiss National Science Foundation (SNSF, Project 315230_130764/1), and by the Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco (FACEPE) and the Universidade Federal de Pernambuco (UFPE) which provided grants to F. Oehl as ‘Visiting Professor’.

Literature cited

- Błaszczkowski J. 1991. Polish *Glomales*. VIII. *Scutellispora* [sic] *nodosa*, a new species with knobby spores. *Mycologia* 83: 537–542.
- Brundrett M, Melville L, Peterson L. 1994. Practical methods in mycorrhizal research. University of Guelph, Mycologue Publications, Guelph, Ontario.
- Goto BT, Maia LC. 2006. Glomerospores, a new denomination for the spores of *Glomeromycota*, a group molecularly distinct from *Zygomycota*. *Mycotaxon* 96: 129–132.
- Goto BT, Maia LC, Oehl F. 2008. *Ambispora brasiliensis*, a new ornamented species in the arbuscular mycorrhiza forming *Glomeromycetes*. *Mycotaxon* 105: 11–18.
- Krüger M, Walker C, Schüßler A. 2011. *Acaulospora brasiliensis* comb. nov. and *Acaulospora alpina* (*Glomeromycota*) from upland Scotland: morphology, molecular phylogeny and DNA-based detection in roots. *Mycorrhiza* 21(6): 577–587. <http://dx.doi.org/10.1007/s00572-011-0361-7>
- Morton JB, Redecker D. 2001. Two new families of *Glomales*, *Archaeosporaceae* and *Paraglomaceae*, with two new genera *Archaeospora* and *Paraglomus*, based on concordant molecular and morphological characters. *Mycologia* 93: 181–195.

- Oehl F, Sýkorová Z, Redecker D, Wiemken A, Sieverding E. 2006. *Acaulospora alpina*, a new arbuscular mycorrhizal fungal species characteristic for high mountainous and alpine grasslands of the Swiss Alps. *Mycologia* 98: 286–294. <http://dx.doi.org/10.3852/mycologia.98.2.286>
- Oehl F, de Souza FA, Sieverding E. 2008. Revision of *Scutellospora* and description of five new genera and three new families in the arbuscular mycorrhiza-forming *Glomeromycetes*. *Mycotaxon* 106: 311–360.
- Oehl F, Jansa J, de Souza FA, Silva GA, 2010. *Cetraspora helvetica*, a new ornamented species in the *Glomeromycetes* from Swiss agricultural fields. *Mycotaxon* 114: 71–84, <http://dx.doi.org/10.5248/114.71>
- Oehl F, Silva GA, Goto BT, Sieverding E. 2011a. *Glomeromycetes*: three new genera and glomoid species reorganized. *Mycotaxon* 116: 75–120. <http://dx.doi.org/116.75>
- Oehl F, Silva DKA, Maia LC, Sousa NMF de, Vieira HEE, Silva GA. 2011b. *Orbispora* gen. nov., ancestral in the *Scutellosporaceae* (*Glomeromycetes*). *Mycotaxon* 116: 161–169. <http://dx.doi.org/10.5248/116.161>
- Oehl F, Silva GA, Goto BT, Maia LC, Sieverding E. 2011c. *Glomeromycota*: two new classes and a new order. *Mycotaxon* 116: 365–379. <http://dx.doi.org/10.5248/116.365>
- Oehl F, Silva GA, Sánchez-Castro I, Goto BT, Maia LC, Vieira HEE, Barea JM, Sieverding E, Palenzuela J. 2011d. Revision of *Glomeromycetes* with entrophosporoid and glomoid spore formation with three new genera. *Mycotaxon* 117: 297–316. <http://dx.doi.org/10.5248/116.297>
- Oehl F, Sýkorová Z, Błaszczowski J, Sánchez-Castro I, Coyne D, Tchabi A, Lawouin L, Hountondji FCC, Silva GA. 2011e. *Acaulospora sieverdingii*, an ecologically diverse new fungus in the *Glomeromycota*, described from lowland temperate Europe and tropical West Africa. *J. Appl. Bot. Food Qual.* 84: 47–53.
- Palenzuela J, Ferrol N, Boller T, Azcón-Aguilar C, Oehl F. 2008. *Otospora bareai*, a new fungal species in the *Glomeromycetes* from a dolomitic shrub-land in the Natural Park of Sierra de Baza (Granada, Spain). *Mycologia* 100(2): 296–305. <http://dx.doi.org/10.3852/mycologia.100.2.296>
- Palenzuela J, Barea JM, Ferrol N, Azcón-Aguilar C, Oehl F. 2010. *Entrophospora nevadensis*, a new arbuscular mycorrhizal fungus, from Sierra Nevada National Park (southeastern Spain). *Mycologia* 102(3): 624–632. <http://dx.doi.org/10.3852/09-145>
- Palenzuela J, Barea JM, Ferrol N, Oehl F. 2011. *Ambispora granatensis*, a new arbuscular mycorrhizal fungus, associated with *Asparagus officinalis* in Andalucía (Spain). *Mycologia* 103(2): 333–340. <http://dx.doi.org/10.3852/09-146>
- Schenck NC, Spain JL, Sieverding E. 1986. A new sporocarpic species of *Acaulospora* (*Endogonaceae*). *Mycotaxon* 25: 111–117.
- Sieverding E. 1988. Two new species of vesicular arbuscular mycorrhizal fungi in the *Endogonaceae* from tropical highlands. *Angew. Bot.* 62: 373–380.
- Sieverding E, Oehl F. 2006. Revision of *Entrophospora*, and description of *Kuklospora* and *Intraspora*, two new genera in the arbuscular mycorrhizal *Glomeromycetes*. *J. Appl. Bot. Food Qual.* 80: 69–81.
- Spain JL. 2003. Emendation of *Archaeospora* and its type species, *Archaeospora trappei*. *Mycotaxon* 87: 109–112.
- Spain JL, Sieverding E, Oehl F. 2006. *Appendicispora*, a new genus in the arbuscular mycorrhizal-forming *Glomeromycetes*, with a discussion of the genus *Archaeospora*. *Mycotaxon* 97: 163–182.
- Walker C. 2008. *Ambispora* and *Ambisporaceae* resurrected. *Mycol. Res.* 112: 297–298. <http://dx.doi.org/10.1016/j.mycres.2008.02.001>

- Walker C, Reed LE, Sanders FE. 1984. *Acaulospora nicolsonii*, a new endogonaceous species from Great Britain. *Trans Br. mycol Soc.* 83: 360–364. [http://dx.doi.org/10.1016/S0007-1536\(84\)80165-5](http://dx.doi.org/10.1016/S0007-1536(84)80165-5)
- Walker C, Vestberg M, Demircik F, Stockinger H, Saito M, Sawaki H, Nishmura I, Schüßler A. 2007. Molecular phylogeny and new taxa in the *Archaeosporales* (*Glomeromycota*): *Ambispora fennica* gen. sp. nov., *Ambisporaceae* fam. nov., and emendation of *Archaeospora* and *Archaeosporaceae*. *Mycol. Res.* 111: 137–153. <http://dx.doi.org/10.1016/j.mycres.2006.11.008>